

Depressive Symptoms and it correlate among Children with Epilepsy at single-center study in Oman

Hamood Al Kiyumi¹, Salim Al Huseini¹, Hassan Mirza², Naser Al Balushi^{2*}, Khalid Al-Risi¹, Alsalt Altoubi¹, Amira Al Hosni², Talal Al-Mashaikhi³, Rola Al Balushi² and Samir Al-Adawi²

¹Psychiatry Residency Training Program, Oman Medical Specialty Board, Muscat, Oman

²Department of Behavioral Medicine, College of Medicine and Health Sciences, Sultan Qaboos University, Muscat, Oman

³Department of Neurology and Epileptology, Sultan Qaboos University Hospital, Muscat, Oman

Received: 10 August 2020

Accepted: 7 March 2021

**Corresponding author:* naser.albalushi@hotmail.com

DOI 10.5001/omj.2021.112

ABSTRACT

Objectives: Epilepsy is the most common neurological disorder in children and comorbid depression is common. This study aimed to assess the frequency of depressive symptoms along with demographic and clinical factors in children diagnosed with epilepsy in a tertiary care institution in Oman.

Methods: This cross-sectional study was conducted between June 2016 and August 2018 and constituted children (n=75) with age group from 6 to 12 years old, attending the epilepsy clinic at a tertiary care center during the study period. The cohort constituted of children with epilepsy (CWE) following up at a dedicated unit. The instrument, Center for Epidemiologic Studies Depression Scale for Children (CES-DC), was utilized to assess the presence of depressive symptoms. Associated factors including history of seizure in the last three months, compliance with antiepileptic medications, and type of epilepsy were also examined.

Results: Depressive symptoms were endorsed in 52% of CWE and 96% of them were compliant to medications. The recurrent seizure was present in the last three months among 48% of the CWE. The type of epilepsy was significantly associated with the presence of depressive symptoms.

Conclusions:The present study found that depression is prevalent among Omani CWE. Certain clinical factors appeared to increase the risk of depression among this population. The findings of this study fill a gap in the existing literature and call for further work aiming to explore possible tailored recognition and CWE.

Keywords: Depressive Symptoms, Children with Epilepsy, neurology clinic,Oman,

Introduction

Epilepsy is the most common neurological disorder in children affecting approximately 41-187/100,000[1]. Children with epilepsy (CWE) exhibit a wide range of psychosocial deficits. Depression is the most prevalent psychiatric manifestation in CWE varying from 10 to 30% [2,3]. A meta-analysis revealed an overall prevalence of 13.5% [4] which exceeds that typically found in children. Depressive symptoms may lead to poor seizure control, reduced quality of life, and suicide [5,6]. Examining epilepsy-specific risk factors revealed that patients who are adolescents have early-onset epilepsy, take two or more antiepileptic medication (AED), and have frequent seizures show an increased risk of developing depressive symptoms [6].As a result, the UK National Institute for Health and Clinical Excellence (NICE) guidelines recommend that the psychological needs of children with epilepsy should be considered as part of routine care [7].The relationship between epilepsy and depression is complex and potentially bidirectional, thereby suggesting common underlying pathophysiology between these conditions[8-11].While rates of epilepsy appear to be waning in high-income countries, the reverse trend appears to be common in the Eastern Mediterranean Region (EMR) [12].A study compared CWE in Northern Jordan with normal children and found that depressive symptoms were observed in approximately 23% of CWE compared to 11% of children without epilepsy [13].As the magnitude of epilepsy tends to strongly hinge on ethnicity [14], there is a dearth of studies examining comorbid depressive symptoms among CWE in the EMR.Due to this,our study explored the frequency of depressive symptoms along with its correlates in children diagnosed with epilepsy in a tertiary care institution in Oman.The covariates in the variation of depressive symptoms were also examined. It is hypothesized that the frequency of depressive symptoms in CWE is higher than in other regions. Additionally, it is expected that sociodemographic factors such as gender and epilepsy-specific risk factors such as the

amount of AED taken will be highly associated with depressive symptoms. This will likely lay the groundwork for the routine and periodic screening of depressive symptoms among CWE for early detection and appropriate management.

Method

Study design and Setting

This is a cross-sectional analytic study that was conducted between June 2016 and August 2018 at the Pediatric Neurology Clinic, Sultan Qaboos University Hospital. Oman has a universal free healthcare system for all citizens and its healthcare structure is divided into primary, secondary, and tertiary care. In Oman, epilepsy clinics are available across the provinces of the country.

Data collection and Participants

The data collection took place while waiting for their appointment to see their doctor in the epilepsy clinic at Sultan Qaboos University Hospital. This study was carried out between June 2016 to August 2018 and included 75 CWE ranging from 6 to 12 years old. Patients were excluded if they were diagnosed with an intellectual disability based on Diagnostic and Statistical Manual of Mental Disorders (Fifth Edition) (DSM-V) criteria [15], already known to have mental illnesses, age below 6 years, those who decline to participate and the forms with missing data were excluded from the study. The caregivers were informed that their children's participation is voluntary and would not affect their clinical care. They were also informed of their right to withdraw at any time. To ensure privacy and avoid feeling coerced to participate, the principal investigator and co-investigators explained the purpose of the study to the participants and the consent to participate was obtained in a private room.

Outcome measures

The Center for Epidemiologic Studies Depression Scale for Children (CES-DC)

The Center for Epidemiologic Studies Depression Scale for Children (CES-DC) [16], was used to measure depressive symptoms. CES-DC is a validated and widely used scale to assess the level of depressive symptoms among children from different clinical populations and across many languages [17,18]. The CES-DC is a 20 item self-report inventory that measures the intensity of depressive symptoms during the past week. It is used for children between ages (6 -12 years old). Children who

struggled to read and understand the questionnaire were assisted by their parents. The questionnaire is scored between 0-60 with a cut-off score of ≥ 15 being indicative of depression [16]. To date, there is not publicly available Arabic version of CES-DC. Therefore, it was necessary to translate it into Arabic. The standard protocol for back translation was utilized as described elsewhere [19]. The final Arabic-version of CES-DC had adequate internal consistency (Cronbach's $\alpha = 0.8$).

Sociodemographic and clinical variable

In addition to CES-DC, various covariates were sought from the accompanying caregivers including gender (male, female), age group (6 to 8 years, 9-12 years). Medical records were screened to document the following clinical risk factors such as seizure history in the last three months (Seizure absent, Seizure present) compliance to medication (Compliant, on-Compliant) according to the parent report and doctor's documentation, and type of epilepsy according to the EEG record (Generalized, focal, Unclassified) were also obtained.

Ethical consideration

This study adheres to the American Association for Public Opinion Research (AAPOR) reporting guidelines [20]. The study was vetted by the local IRB, Medical Research, and Ethical committee at the College of Medicine, Sultan Qaboos University (Ref. No. SQU-EC/148/16, MREC # 1337).

Statistical analysis

The statistical analysis was carried out based on the description, distribution, and categorization of variables included in the study. The significant differences between the presence or absence of depressive symptoms groups for various categorical variables were explored by univariate analysis using the chi-square test. The analyses were performed using IBM SPSS Statistics for Windows version 23.0 [21]. A p-value of less than 0.05 (2-tailed tests) was considered statistically significant.

Results

Table 1 reveals the sociodemographic profile of CWE (n=75) with ages ranging from 6 to 12 years. The study included 45 males (60%) and 30 females (40%) CWE. 96% of CWE were compliant with treatment. 52% (n= 39) CWE reported no seizure in the

past 3 months while 48% (n= 36) CWE reported having at least one seizure episode in the last 3 months.

Regarding the type of epilepsy, 34 participants (45.3%) were having generalized epilepsy, 32 participants (42.7%) were having localized epilepsy, and 12% (n=9) CWE with unclassified epilepsy. There was a significant association between the type of epilepsy and depressive symptoms. Furthermore, children with localized epilepsy reported higher depressive symptoms. The study found that 52% (n=39) CWE endorsed the presence of depressive symptoms. However, there was no association between the presence/absence of depressive symptoms with age, gender, and presence/absence of seizure in the last 3 months as shown in Table 1.

DISCUSSION

To our knowledge, this is the first study that examined the prevalence of depressive symptoms and covariates among children with epilepsy (CWE) in Oman. The main findings suggest that 52% of the sample endorsed depressive symptoms. One way to test the veracity of depressive symptoms is to compare them with other typically developing children. Baki et al. [22] in Istanbul Turkey have reported that CWE attending a tertiary-care setting have higher scores of depressive symptoms using Kovac Child Depression Inventory compared with typically developing children. Loney et al. [23] have employed the Children's Depression Inventory among children presenting with a first seizure in Alberta, Canada. The rate of depressive symptoms was 20% among the cohort. The rate of depressive symptoms appears to be independent of the age of the child [24,25]. Rosic et al. [26] have conducted a prospective longitudinal cohort study of CWE at six Canadian tertiary-care centers. The authors followed those children for over 28 months with repeated assessments using the Children's Depression Inventory Short-Form. The picture that emerged was stark; depressive symptoms remained persistent and pervasive and independent of age, types, and severity of the seizure. Although there is no study on the outcome of resective surgery among CWE, at least in the adult population, resective surgery initially appears to have kickstarted improvement in depressive symptoms, the improvement after resective surgery was more robust among those with superior seizure control.

The prevalence of depressive symptoms among children tends to vary according to the case-ascertainment and measures used to solicit the presence of

depressive symptoms. The presently found 52% of depressive symptoms appear to be in the higher range from previous literature [27]. More studies are therefore warranted to examine the factors that contribute to the wider disparity of the rate of depressive symptoms among CWE.

The second aim of this study was to explore the link between covariates and depressive symptoms, the results revealed a significant association between the type of epilepsy and variation in depressive symptoms. This suggests that children with localized epilepsy are more likely to experience depressive symptoms. This finding echoes previous research that found strong correlations between focal seizures and comorbid psychiatric disorders, specifically depression [3,28,29] but there are dissent views [30]. The findings revealed that children ranging from 9-12 years reported more depressive symptoms and females experienced fewer symptoms of depression, however, the associations were insignificant. The present finding that the female gender has lower depressive symptoms appears to contradict previous studies where depressive symptoms are common in females in the general population as well as those with chronic conditions [31,32]. Future studies should examine whether there are socio-cultural factors that contribute to the present finding. More studies are therefore warranted.

It is expected that younger children report fewer depressive symptoms because younger are not yet capable to introspect and verbalizes emotional state. For example, depressed children fail to express their sadness, instead, their symptoms may appear as anger, irritability, and more importantly changes in their sleeping and eating patterns [33]. Our findings are in the consensus of what has been found in previous studies that females and older age groups with epilepsy were more likely to have depression [3,34]. Moreover, studies were done among children and adolescents attending pediatric outpatient departments of tertiary hospitals found that emotional disorders were 15% among the younger group [35]. Alternatively, parents may be unaware of their children's depressive nature and may underreport such symptoms. This could explain the non-significant association as younger participants were assisted by their parents when completing the questionnaire. Because children with depression are often undiagnosed and untreated, it is important to raise awareness and discuss the likelihood of experiencing depression within CWE. Eastern society assumes that children are mature enough to be afflicted with emotional disorders; this has generally hampered the examination of mood disorders in CWE

[7]. Instead, the presence of internalization behavior problems is attributed to immaturity, sibling rivalry, or simply attention-seeking rather than psychiatric disorders. This may explain the lacking data on mood disorders among CWE in the MER. The magnitude of depressive symptoms in referral hospitals is higher than the global range 5.2% to 39.6% [36]. Asadi-Pooya et al. [37] reviewed studies in Asia that indicated that a high prevalence of depression exists across the continent. This review is limited by the fact that it did not focus on children nor adolescents. Within the EMR, Al-Khateeb & Al-Khateeb [38] also reported similar findings, however, the study was limited to the adult population. Therefore, our study intended to fill the gap in the literature by studying children with epilepsy and examining comorbid depressive symptoms among CWE.

Limitations

Firstly, this study did not include all CWE, only included children up to 12 years rather than up to 18 years, and was hospital-based. Secondly, the scale for screening depression was without the gold-standard face-to-face clinical assessment. Thirdly, AEDs are known to induce dysphoria and negativistic behavior and our study did not consider the relationship between AED and depressive symptoms. Finally, depressive symptoms are strongly associated with the period between seizures -ictal, peri-ictal interictal [39]. Future studies should include studies that control the period between seizures and the types of AED.

Conclusion

Despite the aforementioned limitations, the present study showed that almost half of the children with epilepsy have depressive symptoms. The present study suggests that focal epilepsies are associated with an increased risk of depressive symptoms. Our results suggest that early identification, appropriate referral to child psychiatry for subsequent management are needed. Despite the small sample size, this study is one of the few studies that explored comorbid depression in children with epilepsy in Oman.

Competing / Conflicts of Interest

The authors declare that there are no conflicts of / or competing interests.

Funding

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Acknowledgments

The authors thank the study staff and the staff at Sultan Qaboos University Hospital for their support.

References

1. Camfield P, Camfield C. Incidence, prevalence and aetiology of seizures and epilepsy in children. *Epileptic Disord.* 2015 Jun;17(2):117-23. doi: 10.1684/epd.2015.0736. PMID: 25895502.
2. Adewuya AO, Ola BA. Prevalence of and risk factors for anxiety and depressive disorders in Nigerian adolescents with epilepsy. *Epilepsy Behav.* 2005 May 1;6(3):342–7
3. Caplan R, Siddarth P, Gurbani S, Hanson R, Sankar R, Shields WD. Depression and anxiety disorders in pediatric epilepsy. *Epilepsia.* 2005 May;46(5):720–30
4. Scott AJ, Sharpe L, Loomes M, Gandy M. Systematic review and meta-analysis of anxiety and depression in youth with epilepsy. *Journal of pediatric psychology.* 2020 Jan 6.
5. Johnson EK, Jones JE, Seidenberg M, Hermann BP. The relative impact of anxiety, depression, and clinical seizure features on health-related quality of life in epilepsy. *Epilepsia.* 2004 May;45(5):544-50.
6. Falcone, T., & Timmons-Mitchelle, J. (2013). Pediatric Epilepsy and Depression. Cleveland Clinic. Retrieved from <https://my.clevelandclinic.org/-/scassets/files/org/neurological/epilepsy/pediatric-epilepsy-and-depression-fact-sheet.ashx?la=en>
- 7- NICE: Epilepsies: diagnosis and management (CG137).2012. Reference Source [Google Scholar]
8. Hesdorffer DC, Ishihara L, Mynepalli L, Webb DJ, Weil J, Hauser WA. Epilepsy, suicidality, and psychiatric disorders: a bidirectional association. *Ann Neurol.* 2012;72(2):184–191. doi:10.1002/ana.23601
9. Hesdorffer D. Comorbidity between neurological illness and psychiatric disorders. *CNS Spectr.* 2016;21(3):230–238. doi:10.1017/S1092852915000929

10. Josephson CB, Lowerison M, Vallerand I, et al. Association of depression and treated depression with epilepsy and seizure outcomes: a multicohort analysis. *JAMA Neurol.* 2017;74(5):533–539.
11. Martin R, Faught E, Richman J, et al. Psychiatric and neurologic risk factors for incident cases of new-onset epilepsy in older adults: data from U.S. medicare beneficiaries. *Epilepsia.* 2014;55(7):1120–1127. doi:10.1111/epi.12649
12. Epilepsy in the WHO Eastern Mediterranean Region: Bridging the gap. Regional Office for the Eastern Mediterranean (2010). <https://applications.emro.who.int/dsaf/dsa1039>.
13. Alwash RH, Hussein MJ, Matloub FF. Symptoms of anxiety and depression among adolescents with seizures in Irbid, Northern Jordan. *Seizure.* 2000 Sep 1;9(6):412-6.
14. Hunter MB, Yoong M, Sumpter RE, Verity K, Shetty J, McLellan A, Chin RF. Incidence of early-onset epilepsy: A prospective population-based study. *Seizure.* 2020 Feb 1;75:49-54.
- 15- *Diagnostic and Statistical Manual of Mental Disorders.* 5th. Arlington, VA, USA: American Psychiatric Association; 2013.
16. Weissman MM, Orvaschel H, Padian N: Children's symptom and social functioning self-report scales: Comparison of mothers' and children's reports. *Journal of Nervous Mental Disorders.* 168(12):736–740; 1980.
17. Betancourt T, Scorza P, Meyers-Ohki S, Mushashi C, Kayiteshonga Y, Binagwaho A, Stulac S, Beardslee WR. Validating the center for epidemiological studies depression scale for children in Rwanda. *Journal of the American Academy of Child & Adolescent Psychiatry.* 2012 Dec 1;51(12):1284-92.
18. Essau CA, Olaya B, Pasha G, Gilvarry C, Bray D. Depressive symptoms among children and adolescents in Iran: A confirmatory factor analytic study of the Centre for Epidemiological Studies Depression Scale for Children. *Child Psychiatry & Human Development.* 2013 Feb 1;44(1):123-36.
19. Brislin, R. W. (1980). Translation and content analysis of oral and written materials. In H. C. Triandis & J. W. Berry (Eds.), *Handbook of cross-cultural psychology* (Vol. 2, pp. 389-444). Boston: Allyn & Bacon

20. American Association for Public Opinion Research
<https://www.aapor.org/Publications-Media/AAPOR-Journals/Standard-Definitions.aspx>
21. IBM Corp. Released 2015. IBM SPSS Statistics for Windows, Version 23.0. Armonk, NY: IBM Corp
22. Baki O, Erdogan A, Kantarci O, Akisik G, Kayaalp L, Yalcinkaya C. Anxiety and depression in children with epilepsy and their mothers. *Epilepsy Behav.* 2004 Dec;5(6):958-64. doi: 10.1016/j.yebeh.2004.08.016. PMID: 15582845.
23. Loney, J. C., Wirrell, E. C., Sherman, E. M., & Hamiwka, L. D. (2008). Anxiety and depressive symptoms in children presenting with a first seizure. *Pediatric neurology*, 39(4), 236-240.
24. Ettinger AB, Weisbrot DM, Nolan EE, Gadow KD, Vitale SA, Andriola MR, Lenn NJ, Novak GP, Hermann BP. Symptoms of depression and anxiety in pediatric epilepsy patients. *Epilepsia.* 1998 Jun;39(6):595-9. doi: 10.1111/j.1528-1157.1998.tb01427.x. PMID: 9637601.
25. Dunn DW, Austin JK, Huster GA. Symptoms of depression in adolescents with epilepsy. *J Am Acad Child Adolesc Psychiatry.* 1999 Sep;38(9):1132-8. doi: 10.1097/00004583-199909000-00017. PMID: 10504812.
26. Rosic T, Avery L, Streiner DL, Ferro MA, Rosenbaum P, Cunningham C, Ronen GM. Longitudinal trajectories of depression symptoms in children with epilepsy. *Dev Med Child Neurol.* 2020 May;62(5):593-599. doi: 10.1111/dmcn.14387. Epub 2019 Nov 7. PMID: 31696940.
27. Alamri Y, Al-Busaidi IS. Anxiety and depression in Saudi patients with epilepsy. *Epilepsy Behav.* 2015 Dec 1;53:25.
28. Thome-Souza S, Kuczynski E, Assumpção F Jr, et al. Which factors may play a pivotalrole on determining the type of psychiatric disorder in children and adolescents withepilepsy?. *Epilepsy Behav.* 2004;5(6):988-994.
29. Jones JE, Watson R, Sheth R, et al. Psychiatric comorbidity in children with new onset epilepsy. *Dev Med Child Neurol.* 2007;49(7):493-497. doi:10.1111/j.1469-8749.2007.00493.x
30. Rosic T, Avery L, Streiner DL, Ferro MA, Rosenbaum P, Cunningham C, Ronen GM. Longitudinal trajectories of depression symptoms in children with epilepsy. *Dev Med Child Neurol.* 2020 May;62(5):593-599. doi: 10.1111/dmcn.14387. Epub 2019 Nov 7. PMID: 31696940.

31. Piquart M, Shen Y. Depressive symptoms in children and adolescents with chronic physical illness: an updated meta-analysis. *J Pediatr Psychol*. 2011 May;36(4):375-84. doi: 10.1093/jpepsy/jsq104. Epub 2010 Nov 18. PMID: 21088072.
32. Gaus V, Kiep H, Holtkamp M, Burkert S, Kendel F. Gender differences in depression, but not in anxiety in people with epilepsy. *Seizure*. 2015 Nov 1;32:37-42.
33. Cole PM, Luby J, Sullivan MW. Emotions and the Development of Childhood Depression: Bridging the Gap. *Child Dev Perspect*. 2008;2(3):141-148.
34. Oguz A, Kurul S, Dirik E. Relationship of epilepsy-related factors to anxiety and depression scores in epileptic children. *J Child Neurol*. 2002;17(1):37-40.
35. Jesmin A; Mullick MSI, Rahman KM, Muntasir MM. Psychiatric Disorders in Children and Adolescents Attending Pediatric Out Patient Departments of Tertiary Hospitals. *Oman Med J*. 2016;31(4):258-262. doi:10.5001/omj.2016.51
36. Maryam S, Parviz B. Depression in children and adolescents with epilepsy: a 15 year research review of prevalence, and demographic and seizure related correlates. *Iran J Pediatr*. 2013;23(1):1-7.
37. Asadi-Pooya AA, Kanemoto K, Kwon OY, Taniguchi G, Dong Z, Chinvarun Y, Yu HY, McGonigal A, Kanner AM, Park SP. Depression in people with epilepsy: How much do Asian colleagues acknowledge it?. *Seizure*. 2018 Apr 1;57:45-9.
38. Al-Khateeb JM, Al-Khateeb AJ. Research on psychosocial aspects of epilepsy in Arab countries: a review of literature. *Epilepsy & behavior*. 2014 Feb 1;31:256-62.
39. Kanner AM. Depression in epilepsy: prevalence, clinical semiology, pathogenic mechanisms, and treatment. *Biol Psychiatry* 2003; 54: 388–98.

Table1: Sociodemographic profile and Association of depressive symptoms within specific categories among 75 children diagnosed with epilepsy seeking consultation at the Pediatric Neurology Clinic.

Variable	N (%)	Depressive symptoms		Chi-square p-value	
		Present (n)	Absent (n)		
		39 (52%)	36 (48%)		
Age group	6 to 8 years	29 (38.7)	14	15	NS
	9 to 12 years	46 (61.3)	25	21	
Gender	Male	45 (60)	26	19	NS
	Female	30 (40)	13	17	
Seizure in the last 3 months	Present	36 (48)	22	14	0.09
	Absent	39 (52)	17	22	
Type of Epilepsy	Generalized	34 (45.3)	10	24	0.002
	focal	32 (42.7)	23	9	

	Unclassified	9 (12)	6	3	
Compliance to medications	Compliant	72 (96)	39	33	0.106
	Non-compliant	3 (4)	0	3	