

# Evaluation of *in Vitro* Activity of Ceftazidime/Avibactam and Ceftolozane/Tazobactam against ESBL-producing Enterobacterales Isolated from Intensive Care Units from Qatar

Mazen A. Sid Ahmed<sup>1,2\*</sup>, Emad Bashir Ibrahim<sup>1,3</sup>, Jemal M. Hamid<sup>1</sup>, Joanne Daghfal<sup>4</sup>, Mohammed A. Alyazidi<sup>1</sup>, Aimen H. Abdelwahab<sup>1</sup>, Muna A. Al-Maslamani<sup>4,5</sup>, Abdul Latif Al Khal<sup>4,5</sup> and Hamad Abdel Hadi<sup>4,5</sup>

<sup>1</sup>Division of Microbiology, Department of Pathology and Laboratory Medicine, Hamad Medical Corporation, Doha, Qatar

<sup>2</sup>The Life Science Centre, School of Science and Technology, Örebro University, Örebro, Sweden

<sup>3</sup>Biomedical Research Centre, Qatar University, Doha, Qatar

<sup>4</sup>Communicable Diseases Center, Hamad Medical Corporation, Doha, Qatar

<sup>5</sup>Division of Infectious Diseases, Department of Medicine, Hamad Medical Corporation, Doha, Qatar

## ARTICLE INFO

### Article history:

Received: 27 December 2021

Accepted: 12 April 2022

### Online:

DOI 10.5001/omj.2022.89

### Keywords:

Enterobacterial; beta-lactamase TEM-11; Antimicrobial drug resistance; Ceftazidime-Avibactam; Ceftolozane-Tazobactam; Qatar.

## ABSTRACT

**Objectives:** Extended-spectrum  $\beta$ -lactamases (ESBLs) mechanism of resistance in Enterobacterales leads to poor clinical outcomes. Ceftazidime/avibactam and ceftolozane/tazobactam are two broad-spectrum antimicrobial combinations that are effective against multidrug-resistant organisms with regional variations. This study aims to evaluate the antimicrobial susceptibility test (AST) for both combinations against ESBL-producing Enterobacterales isolated from intensive care units (ICUs) in tertiary hospitals from November 2012 to October 2013 in Qatar. **Methods:** A total of 629 Enterobacterales isolates from ICUs were screened for ESBL production using BD-Phoenix™ confirmed by double-disk potentiation, while ESBL-genes were detected by polymerase chain reaction. The ASTs for ceftazidime/avibactam and ceftolozane/tazobactam were assessed by minimum inhibitory concentration (MIC) test strips. A single isolate that was resistant to both combinations was subjected to whole-genome sequencing. **Results:** The prevalence of ESBL-producing Enterobacterales isolated from ICUs was 17.3% (109/629) with predominance of *Klebsiella pneumoniae* (56/109; 51.4%) and *Escherichia coli* (38/109; 34.9%). The susceptibility of ceftazidime/avibactam and ceftolozane/tazobactam against ESBL-producers was 99.1% (108/109) and most (81/109; 74.3%) had MICs < 0.5 for both combinations. The predominant ESBL-gene was *bla*<sub>CTX-M</sub> (72/109; 66.1%). A single isolate that was resistant to both combinations harbored multiple ESBL resistant-genes including *bla*<sub>VEB-5</sub> and *bla*<sub>VIM-2</sub>. **Conclusions:** ESBL-producing Enterobacterales isolated from ICUs were predominantly *K. pneumoniae* and *E. coli*, mainly harboring *bla*<sub>CTX-M</sub> gene. They were highly susceptible to ceftazidime/avibactam and ceftolozane/tazobactam suggesting potential alternatives to currently available therapeutic options.

The management of infections secondary to multidrug-resistant organisms (MDROs) that encompasses gram-negative bacteria (GNB) is a global healthcare challenge not only because of the limited available treatment options but also for their associated significant morbidity and mortality as well as the substantial cost of management.<sup>1,2</sup>

In secondary and tertiary hospitals, the ultimate antimicrobial resistance (AMR) is encountered at intensive care units (ICUs) where the critical nature

of patients' cohort, concurrent comorbidities, invasive procedures, prior colonization as well as environmental exposure to MDROs that is accelerated by high antibiotics consumption are inevitable acquisition hazards.<sup>3,4</sup> Over the past decade in Qatar, internal microbiological surveillance and monitoring of GNB, particularly the Enterobacterales, has revealed alarmingly rising trends of AMR, particularly for extended-spectrum  $\beta$ -lactamases (ESBLs) in line with shifting regional epidemiology.<sup>5,6</sup> In critical care settings typical

recommended approach for the management of ESBL-producing Enterobacterales is treatment with carbapenems, particularly if there is an associated serious invasive or high inoculum disease.<sup>7</sup> The concern of diminishing efficacy of the limited treatment options against the ever-rising resistant bacterial strains, led infection specialists to seek alternatives to carbapenem therapy.

Ceftazidime/avibactam and ceftolozane/tazobactam are  $\beta$ -lactam/ $\beta$ -lactamase inhibitors (BLBLIs) combinations that are approved by both the United States Food and Drug Administration and the European Medicines Agency, demonstrating comparable or superior activity against MDROs particularly in GNB for the treatment of complicated urinary tract and intra-abdominal infections as well as infections secondary to hospital or ventilation associated pneumonia.<sup>8</sup> Avibactam is a non-BLBLI that potently inhibits most (but not all) class A ESBLs, class C (including AmpC enzymes), and some class D  $\beta$ -lactamases.<sup>9</sup> Furthermore, due to its different mode of action, avibactam is considered as one of the most effective BLBLIs displaying a broader inhibitory range and spectrum.<sup>10</sup> On the other hand, ceftolozane is a novel cephalosporin that is not affected by outer membrane protein loss which is a weak substrate for drug efflux pump mechanism, rendering the drug exhibiting less affinity for hydrolysis by AmpC, and hence better efficacy.<sup>11</sup> Pairing ceftolozane with the classic  $\beta$ -lactamase inhibitor tazobactam has broadened their capacity to act on most ESBL-producing GNB.<sup>12</sup>

The presented study aims mainly to evaluate the antimicrobial activity of ceftazidime/avibactam and ceftolozane/tazobactam against 109 ESBL-producing Enterobacterales isolates from ICUs in Qatar,<sup>13</sup> describe its microbiological characteristics, as well as the underlying genomic resistance profiles.

## METHODS


This research project was approved by the Institutional Review Board at Hamad Medical Corporation (HMC), which complies with international ethical standards and regulations (Protocol no. RC/75813/2013). The study was conducted on routine specimens processed by the Microbiology Division, Department of Laboratory Medicine and Pathology, HMC, Qatar. All samples were collected prospectively over one year (1 November 2012 to 31

October 2013) from patients admitted to all ICUs (medical 29%, surgical 29%, trauma 16%, pediatric 16%, and neonatal 10%) at HMC. These were then analyzed for the presence of resistant pathogens.

The study definitions recognized duplicates of the same species of bacteria as isolates from the same patient displaying identical antimicrobial susceptibility patterns when isolated within 30 days regardless of sample sites which were considered repetitive and excluded. Isolates with major differences in antimicrobial susceptibilities were counted as new even within the defined 30 days time frame. The single isolate that was resistant to ceftazidime/avibactam and ceftolozane/tazobactam underwent standard diagnostic work-up, and then was stored at  $-80^{\circ}\text{C}$  pending further genomic analysis.

Microbiological identification and antimicrobial susceptibility tests (AST) were performed using BD Phoenix<sup>TM</sup> automated system according to manufacturer recommendations. Samples that tested positive for ESBL by Phoenix, or showed a minimum inhibitory concentration (MIC) of  $> 8 \mu\text{g}/\text{mL}$  for 3<sup>rd</sup> generation cephalosporins or aztreonam, were subsequently confirmed by a double-disk potentiation test with ceftazidime, amoxicillin/clavulanic acid, ceftriaxone, and ceftoxitin antibiotics, interpreted as recommended by Clinical Laboratory Standards Institute standards for ESBL identification.<sup>14</sup> AST and MIC for ceftazidime/avibactam and ceftolozane/tazobactam were performed using MIC Test Strips (Liofilchem<sup>®</sup>, Diagnostics, Italy). *Escherichia coli* ATCC 25922, *E. coli* ATCC 35218, and *Pseudomonas aeruginosa* ATCC 27853 were used as controls. Susceptibility reporting was based on the Clinical Laboratory Standards Institute recommendations.<sup>14</sup> Since there were no recommended intermediate susceptibility categories available for ceftazidime/avibactam against Enterobacterales, isolates were described as 'susceptible' if the MIC was  $\leq 8 \text{ mg}/\text{L}$  and 'non-susceptible' if the MIC was  $> 8 \text{ mg}/\text{L}$  as outlined in the supplementary Table A1 in Appendix.<sup>14</sup> To achieve consistency, intermediate and resistant categories were grouped as non-susceptible for all reported antimicrobial agents.

Bacterial DNA extraction and detection of ESBL resistance genes were performed through an in-house polymerase chain reaction (PCR) technique, using the boiling lysis methods.<sup>15</sup> PCR reactions for the



**Figure 1:** Antimicrobial susceptibility results for ceftazidime/avibactam, ceftolozane/tazobactam, and comparator agents against clinical extended-spectrum  $\beta$ -lactamase-producing Enterobacteriales isolates from Qatar.

ESBL genes (TEM, SHV, and CTX-M-1) were conducted using previously described protocols.<sup>16</sup>

Whole-genome sequencing (WGS) was performed to study isolated genomic relationships for annotating antibiotic resistance genes (ARGs). Extracted DNA was sent to GATC Service (Eurofins Genomics, Germany) for sequencing using Illumina HiSeq 2000 system (Illumina, San Diego, California). The genes were assembled using SPAdes, Version 3.13.0 (<https://cab.spbu.ru/software/spades/>) while Multi-locus sequence typing (MLST) of the described resistant isolate of *E. coli* was performed on MLST server 1.8 provided (<https://cge.cbs.dtu.dk/services/MLST/>). ARGs were annotated using Comprehensive Antibiotic Resistance Database (CARD), Version 1.2.0 (<https://card.mcmaster.ca/>).

Demographics of patients, characteristics of isolates, as well as the patterns of antimicrobial susceptibility of ESBL-producing Enterobacteriales including resistant genes were presented as numbers and percentages using Stata statistical software (Stata Corp LLC, College Station, Texas version 16.1).

## RESULTS

Out of 629 Enterobacteriales isolates investigated, 109 (17.3%) isolates from 87 patients were found to be ESBL positive. The source samples of these were:

respiratory 35.8% (39/109), blood 27.5% (30/109), urine 24.8% (27/109), fluids 6.4% (7/109), and others 5.5% (6/109). The ESBL-positive isolates were predominantly *Klebsiella pneumoniae* (50.5%) and *E. coli* (34.9%) while other species comprised 13.7%. The majority of isolates were from male patients 65 (59.6%) and those aged between one month and 86 years. The patients were categorized into three age groups labeled adult, pediatric, and geriatrics. 'Adult' group (14–65 years) contributed to more than half (57/109; 52.3%) of the resistant isolates, followed by 'pediatric' (9/109; 8.3%) < 14 years, and 'geriatric' (26/109; 23.9%) > 65 years.

The predominantly identified ESBL-producing genes were *bla*<sub>CTX-M-1</sub> (72/109; 66.1%) followed by *bla*<sub>SHV</sub> (58/109; 53.2%) and *bla*<sub>TEM</sub> (44/109; 40.4%). All three  $\beta$ -lactamase genes (TEM, SHV, and CTX-M-1) were detected in 26/56 (46.4%) of *K. pneumoniae* isolates, while two genes (SHV/CTX-M-1) were present in 10/56 (17.8%) of *K. pneumoniae* and only 1/38 (2.6%) of *E. coli* isolates, with TEM/CTX-M-1 being present in 7/38 (18.4%) of *E. coli* and 4/56 (7.1%) of *K. pneumoniae*, while TEM/SHV was detected in only 2/38 (5.3%) of *E. coli* isolates.

The activity of ceftazidime/avibactam and ceftolozane/tazobactam against 109 ESBL-producing Enterobacteriales isolates demonstrated 99.1% (108/109) susceptibility for both combinations. Only meropenem showed 100% (109/109) susceptibility followed by imipenem at 99.1% while ertapenem and amikacin susceptibility was 97.2%. Other antimicrobials demonstrated moderate-to-low susceptibility rates with 78.0% for piperacillin/tazobactam, 64.2% for tigecycline, 60.6% for ciprofloxacin, and 38.5% for cotrimoxazole while as predicted cephalosporin had high-level resistance (99.1% for ceftriaxone and 93.6% for cefepime) [Figure 1]. Furthermore, most of the ESBL-producing Enterobacteriales were highly susceptible to ceftazidime/avibactam at low MICs (MIC<sub>50/90</sub> 0.19/0.38  $\mu$ g/mL) and ceftolozane/tazobactam (MIC<sub>50/90</sub> 0.38/1.00  $\mu$ g/mL) [Table 1], with the majority of isolates demonstrating MICs < 0.5 (n = 81; 74.3%) [Table 2]. The additional microbiological and molecular characterization including susceptibility testing results are shown in Appendix [Table A1].

Our findings are distinctively different from other regional studies where ceftazidime/avibactam

**Table 1:** Minimum inhibitory concentration (MIC) for ceftazidime/avibactam and ceftolozane/tazobactam against 109 clinical extended-spectrum  $\beta$ -lactamase-producing Enterobacterales isolates collected from intensive care units, Hamad Medical Corporation, Qatar

Organism	Number of isolates	Antibiotic	Range	Susceptible isolates, n (%)	MIC50	MIC90
<i>Klebsiella pneumoniae ssp pneumoniae</i>	55	CZA	0.09–0.75	55 (100)	0.25	0.38
		C/T	0.25–1.50	55 (100)	0.38	1.00
<i>Escherichia coli</i>	38	CZA	0.06–256.00	37 (97.4)	0.12	0.38
		C/T	0.19–256.00	37 (97.4)	0.38	0.75
<i>Enterobacter aerogenes</i>	4	CZA	0.09–0.25	4 (100)	0.19	0.25
		C/T	0.38–0.50	4 (100)	0.38	0.50
<i>Enterobacter cloacae</i>	4	CZA	0.09–0.19	4 (100)	0.09	0.19
		C/T	0.19–0.25	4 (100)	0.25	0.25
<i>Serratia marcescens</i>	2	CZA	0.02–0.12	2 (100)	0.02	0.12
		C/T	0.19	2 (100)	0.19	0.19
<i>Citrobacter braakii</i>	1	CZA	0.25	1 (100)	0.25	0.25
		C/T	0.75	1 (100)	0.75	0.75
<i>Citrobacter freundii</i>	1	CZA	0.06	1 (100)	0.06	0.06
		C/T	0.38	1 (100)	0.38	0.38
<i>Citrobacter amalonaticus</i>	1	CZA	0.12	1 (100)	0.12	0.12
		C/T	0.19	1 (100)	0.19	0.19
<i>Klebsiella oxytoca</i>	1	CZA	0.12	1 (100)	0.12	0.12
		C/T	0.38	1 (100)	0.38	0.38
<i>Klebsiella pneumoniae ssp ozaenae</i>	1	CZA	0.09	1 (100)	0.09	0.09
		C/T	0.25	1 (100)	0.25	0.25
<i>Proteus penneri</i>	1	CZA	0.04	1 (100)	0.04	0.04
		C/T	1.00	1 (100)	1.00	1.00
Total	109	CZA	0.02–256.00	108 (99.1)	0.19	0.38
		C/T	0.19–256.00	108 (99.1)	0.38	1.00

CZA: ceftazidime-avibactam; C/T: ceftolozane-tazobactam.

**Table 2:** Comparison of minimum inhibitory concentration for ceftazidime/avibactam vs ceftolozane/tazobactam against 109 clinical extended-spectrum  $\beta$ -lactamase-producing Enterobacterales isolates from samples from intensive care unit patients at Hamad Medical Corporation, Qatar.

Antibiotic	MIC	Ceftolozane/tazobactam, n (%)					Total
		< 0.25	< 0.5	< 0.75	< 4	> 256	
Ceftazidime/avibactam	< 0.1	13 (11.9)	6 (5.5)	1 (0.9)	1 (0.9)	0.0	21 (19.3)
	< 0.25	7 (6.4)	13 (11.9)	0.0	0.0	0.0	20 (18.3)
	< 0.5	5 (4.6)	37 (33.9)	14 (12.8)	7 (6.4)	0.0	63 (57.8)
	< 0.75	0.0	0.0	0.0	3 (2.8)	0.0	3 (2.8)
	< 1	0.0	1 (0.9)	0.0	0.0	0.0	1 (0.9)
	> 256	0.0	0.0	0.0	0.0	1 (0.9)	1 (0.9)
	Total		25 (22.9)	57 (52.3)	15 (13.8)	11 (10.1)	1 (0.9)

demonstrated superior activity when compared to ceftolozane/tazobactam against ESBL-producer [Table 3], which suggests a potential correlation of embedded ESBL resistance genes not demonstrated in our study because of paucity of resistant isolates [Table 4].<sup>22</sup>

Among the 109 identified ESBL-producing Enterobacterales only one (0.9%) *E. coli* isolate was completely resistant to both ceftolozane/tazobactam and ceftolozane/tazobactam, with MIC > 256 [Table 1]. The resistant isolate was collected from peritoneal fluid of a fatal case of complicated



**Table 3:** Summary of studies comparing *in vitro* activity of ceftazidime/avibactam and ceftolozane/tazobactam against extended-spectrum  $\beta$ -lactamase-producing Enterobacteriales from different geographical regions worldwide.

Study	Geographic location	Susceptibility testing method	Inclusion criteria	Collection years	Number included	Susceptible to MEM, n (%)	Susceptible to CZAn, (%)	Susceptible to C/T, n (%)
Alatoom et al, <sup>17</sup> 2017	Abu Dhabi, UAE	Etest	Resistant to $\geq 1$ agent from $\geq 3$ antimicrobial classes	2015–2016	31	NA	29 (93.5)	30 (96.8)
Sader et al, <sup>18</sup> 2020	70 medical centers, USA	Broth microdilution	ESBL-producing Enterobacteriales from patients hospitalized with pneumonia	2017–2018	285	283 (99.3)	285 (100)	219 (76.8)
Viala et al, <sup>19</sup> 2019	Montpellier, France	Etest	3rd G cephalosporin resistant Enterobacteriaceae	2017	62	NA	60 (97)	34 (65)
Araj et al, <sup>20</sup> 2020	Beirut, Lebanon	MIC gradient Strip Test	MDR and ESBLs <i>E. coli</i> and <i>K. pneumoniae</i>	2017–2018	199	NA	NA	159 (79.9)
Hirsch et al, <sup>21</sup> 2020	Boston, MA; and Philadelphia, PA	Broth microdilution	carbapenem-susceptible (meropenem MIC $\leq 1$ mg/L)	2013–2016	119	119 (100)	119 (100)	109 (91.6)

CZ-A: ceftazidime/avibactam; C/T: ceftolozane/tazobactam; MDR: multi-drug resistant; MEM: meropenem; NS: non-susceptible. \*All studies reported the isolates as susceptible if the MIC was  $\leq 8$  mg/L for ceftazidime/avibactam and  $\leq 4$  mg/L for ceftolozane/tazobactam.

**Table 4:** Genotypic profiles of different  $\beta$ -lactamase enzymes detected among extended-spectrum  $\beta$ -lactamase-producing *E. coli* isolated from samples from intensive care unit patients at Hamad Medical Corporation, Qatar.

Resistance gene	Gene family	Gene identity, %
CTX-M-15	Class A $\beta$ -lactamase	100
VEB-5	Class A $\beta$ -lactamase	100
VIM-2	Class B $\beta$ -lactamase	100
<i>E. coli</i> ampC	Class C $\beta$ -lactamase	97.9
<i>E. coli</i> ampC1	Class C $\beta$ -lactamase	99.3
<i>E. coli</i> ampH	Class C $\beta$ -lactamase	99.2
CMY-42	Class C $\beta$ -lactamase	100
OXA-10	Class D $\beta$ -lactamase	100
OXA-4	Class D $\beta$ -lactamase	100
OXA-486	Class D $\beta$ -lactamase	100

intra-abdominal infection, and was subsequently identified as sequence type ST38. Genomic data analysis revealed that the resistant isolate possessed different ARGs including 11 different  $\beta$ -lactamase genes from all classes; Class A ESBL (CTX-M-1 and VEB-5), Class B metallo- $\beta$ -lactamase (MBL) including *bla*<sub>VIM-2</sub>, class C  $\beta$ -lactamase including *bla*<sub>PCMY-42</sub>. Class D  $\beta$ -lactamase such as *bla*<sub>OXA-4</sub>, *bla*<sub>OXA-10</sub>, and *bla*<sub>OXA-486</sub> [Table 4].

## DISCUSSION

AMR is a major global healthcare challenge with ominous outcomes. Its ultimate manifestation occurs at critical care units where potent risk factors converge—such as a hazardous environment, vulnerable host, and highly resistant pathogens.<sup>23</sup> Thus, one of the foremost challenges in critical care is prevention and management of infections caused by MDR gram-negative organisms, particularly ESBL-producing Enterobacteriales resistant to most antimicrobial classes including most  $\beta$ -lactam penicillins, BLBLIs, and cephalosporins.<sup>23,24</sup> To combat the growing problem of ESBL-producing Enterobacteriales, recent decades have witnessed exponentially rising reliance on carbapenems, to the point of their becoming the *sine qua non* for its management, especially in the context of invasive or high-burden disease.<sup>23,25</sup> In complicated ESBL infections, randomized control trials have demonstrated the superiority of carbapenems over comparators including BLBLIs.<sup>25</sup> However, the near-

universal use of carbapenems has not prevented the problem from increasing; hence, the relentless search for ever-more powerful antibiotic regimens.<sup>8</sup> Among the most promising new combinations are ceftazidime/avibactam and ceftolozane/tazobactam. Several global studies have evaluated the spectrum of their efficacy *in vitro* and *in vivo* against highly resistant gram-negative strains including ESBL producers and their regional variations.<sup>17–21,26</sup>

A 2010 study in Qatar that investigated 450 episodes of invasive bacteremia from a single institution found 61% prevalence of GNB, the most prevalent of which were *E. coli* (27.8%) and *K. pneumoniae* (17.9%), most of them ESBL producers.<sup>27</sup> The scale of the growing problem at the same institution became clearer in a 2020 study which analyzed the results of culture-positive complicated urinary tract infections among adults admitted to surgical ICUs over a 10-year period (2008–2018). The study found that 36% of the isolated pathogens were ESBLs.<sup>28</sup> Healthcare leaders in the Gulf Cooperation Council countries have recognized the problem of resistant pathogens as a health priority and initiated regional collaboration against this common threat.<sup>29</sup>

In our study, 99.1% of ESBL isolates were highly susceptible and most isolates (74.3%) exhibited MIC < 0.5 for both ceftazidime/avibactam and ceftolozane/tazobactam [Table 2]. Notably, the observed high-level susceptibility for ceftazidime/avibactam and ceftolozane/tazobactam against ESBL-producing Enterobacterales isolates collected from prospective critical care clinical cases, predates the introduction of these agents into clinical practice in Qatar. Our microbiological evaluation suggests these novel agents might be rational empirical treatment options sparing carbapenems.

In the Arabian Gulf region, the high volume of international travel coupled with population diversity and high antibiotic consumption are contributing factors towards the rising and diversifying trends of ESBLs in GNB. The problem has reached an endemic state requiring alternative management options.<sup>6,30</sup>

In line with regional and global ESBL genomic studies, the observation that *bla*<sub>CTX-M</sub> in conjunction with *bla*<sub>SHV</sub> and *bla*<sub>TEM</sub> are the main ARGs for ESBL-producing Enterobacterales, points towards the role of cephalosporins as the potential driving precipitant.<sup>5,21,31</sup> In Qatar, the

molecular epidemiology of Enterobacterales from the pediatric population follows the same trends in the region when a large study of 327 sequenced ESBL producers from clinical samples at the largest children's hospital in the region demonstrated dominance of *E. coli* and *K. pneumoniae* as main pathogens with predominance of *bla*<sub>CTX-M-1</sub> and coproduction of *bla*<sub>OXA-1</sub> and *bla*<sub>TEM-1B</sub> as ARGs.<sup>32</sup> In contrast, in the adults population, there are no detailed recent studies to evaluate the wider molecular epidemiology of ESBL in the country but the study of 149 non-repetitive carbapenem-resistant Enterobacterales confirmed regional preponderance of *bla*<sub>NDM</sub> and *bla*<sub>OXA48</sub>.<sup>33</sup>

Not surprisingly, following undergoing WGS, the only concomitant isolate resistant to both ceftazidime/avibactam and ceftolozane/tazobactam harbored multitude of different ARGs.

The ESBL-producing *E. coli* which belonged to ST38 possessed  $\beta$ -lactamase genes from all classes as shown in Table 3. Intriguingly, the detailed study demonstrated the presence of *bla*<sub>VIM-2</sub> MBL which is known to play a fundamental role in ceftazidime/avibactam and ceftolozane/tazobactam resistance.<sup>34</sup> In addition to the endemic class A *bla*<sub>CTX-M</sub> the resistant isolate also harbored *bla*<sub>VEB-5</sub>, which was initially detected in *E. coli* in the USA (GenBank accession number EF420108). The ARG, *bla*<sub>VEB</sub> confers high-level resistance to cephalosporins as well as monobactams and has been shown to inactivate ceftolozane/tazobactam.<sup>35</sup> However, *bla*<sub>VEB-5</sub> is known to be inhibited by avibactam which restored the MIC of ceftazidime from 256  $\mu$ g/mL to 2  $\mu$ g/mL for ceftazidime/avibactam combination.<sup>36</sup> In addition to that, the resistant isolate has multiple underlying ARGs including *bla*<sub>PCMY-42</sub> (AmpC), which drives ceftolozane/tazobactam resistance<sup>37</sup> as well as class D  $\beta$ -lactamases *bla*<sub>OXA-10</sub>, which has been recently reported to enhance ceftolozane/tazobactam and ceftazidime/avibactam resistance.<sup>38</sup>

Despite its wide mechanism of action against MDROs including class A, C, and D  $\beta$ -lactamases, both ceftazidime/avibactam and ceftolozane/tazobactam remain vulnerable when encountering embedded class B  $\beta$ -lactamases such as the potent carbapenemase *bla*<sub>VIM-2</sub> MBL, as described in the isolate of the study.<sup>21,39</sup> Although some molecular tests have been developed to screen for ceftazidime/avibactam and ceftolozane/tazobactam resistance, the current recommendations to interpret activity

through the golden routes of ASTs hold true.<sup>40</sup>

As a consequence, from our study, the prime recommendation is to conduct an urgent clinical evaluation of the novel antibiotics as alternative therapeutic options for MDROs including ESBLs, particularly in critical care settings. This should be supported by data from surveillance and monitoring mechanisms to evaluate the prevalence and characteristics of AMR in the region.

## CONCLUSION

ESBL-producing Enterobacterales represent a significant and growing threat to healthcare in Qatar and the Arabian Gulf region in general, particularly in critical care settings. MDROs such as *K. pneumoniae* and *E. coli* harboring multiple ARGs continue to predominate. Promising high *in vitro* antimicrobial susceptibility to ceftazidime/avibactam and ceftolozane/tazobactam against ESBLs-producing Enterobacterales have added to the arsenal of alternative management options to overcome the growing resistance problem.

### Disclosure

The authors declared no conflicts of interest. The publication of this article was funded by the Qatar National Library. Funders were not involved in the conduct of the study, preparation of the manuscript or the decision to submit the manuscript for publication.

### REFERENCES

- Naylor NR, Atun R, Zhu N, Kulasabanathan K, Silva S, Chatterjee A, et al. Estimating the burden of antimicrobial resistance: a systematic literature review. *Antimicrob Resist Infect Control* 2018 Apr;7:58.
- Wozniak TM, Barnsbee L, Lee XJ, Pacella RE. Using the best available data to estimate the cost of antimicrobial resistance: a systematic review. *Antimicrob Resist Infect Control* 2019 Feb;8:26.
- Massart N, Camus C, Benezit F, Moriconi M, Fillatre P, Le Tulzo Y. Incidence and risk factors for acquired colonization and infection due to extended-spectrum beta-lactamase-producing gram-negative bacilli: a retrospective analysis in three ICUs with low multidrug resistance rate. *Eur J Clin Microbiol Infect Dis* 2020 May;39(5):889-895.
- Teerawattanapong N, Kengkla K, Dilokthornsakul P, Saokaew S, Apisarnthanarak A, Chaiyakunapruk N. Prevention and control of multidrug-resistant gram-negative bacteria in adult intensive care units: a systematic review and network meta-analysis. *Clin Infect Dis* 2017 May;64(suppl\_2):S51-S60.
- Zowawi HM, Balkhy HH, Walsh TR, Paterson DL.  $\beta$ -Lactamase production in key gram-negative pathogen isolates from the Arabian Peninsula. *Clin Microbiol Rev* 2013 Jul;26(3):361-380.
- Dandachi I, Chaddad A, Hanna J, Matta J, Daoud Z. Understanding the epidemiology of multi-drug resistant gram-negative bacilli in the Middle East using a one health approach. *Front Microbiol* 2019 Aug;10:1941.
- Rupp ME, Fey PD. Extended spectrum beta-lactamase (ESBL)-producing Enterobacteriaceae: considerations for diagnosis, prevention and drug treatment. *Drugs* 2003;63(4):353-365.
- van Duin D, Bonomo RA. Ceftazidime/Avibactam and Ceftolozane/Tazobactam: Second-generation  $\beta$ -Lactam/ $\beta$ -Lactamase Inhibitor Combinations. *Clin Infect Dis* 2016 Jul;63(2):234-241.
- Nichols WW, Newell P, Critchley IA, Riccobene T, Das S. Avibactam pharmacokinetic/pharmacodynamic targets. *Antimicrob Agents Chemother* 2018 May;62(6):62.
- Tuon FF, Rocha JL, Formigoni-Pinto MR. Pharmacological aspects and spectrum of action of ceftazidime-avibactam: a systematic review. *Infection* 2018 Apr;46(2):165-181.
- Shorridge D, Pfaller MA, Castanheira M, Flamm RK. Antimicrobial activity of ceftolozane-tazobactam tested against Enterobacteriaceae and *Pseudomonas aeruginosa* collected from patients with bloodstream infections isolated in United States hospitals (2013-2015) as part of the Program to Assess Ceftolozane-Tazobactam Susceptibility (PACTS) surveillance program. *Diagn Microbiol Infect Dis* 2018 Oct;92(2):158-163.
- Sucher AJ, Chahine EB, Cogan P, Fete M. Ceftolozane/tazobactam: a new cephalosporin and  $\beta$ -lactamase inhibitor combination. *Ann Pharmacother* 2015 Sep;49(9):1046-1056.
- Sid Ahmed MA, Bansal D, Acharya A, Elmi AA, Hamid JM, Sid Ahmed AM, et al. Antimicrobial susceptibility and molecular epidemiology of extended-spectrum beta-lactamase-producing Enterobacteriaceae from intensive care units at Hamad Medical Corporation, Qatar. *Antimicrob Resist Infect Control* 2016 Feb;5:4.
- Clinical Laboratory Standards Institute. Performance standards for antimicrobial susceptibility testing. Wayne, PA, USA; 2020.
- Queipo-Ortuño MI, De Dios Colmenero J, Macias M, Bravo MJ, Morata P. Preparation of bacterial DNA template by boiling and effect of immunoglobulin G as an inhibitor in real-time PCR for serum samples from patients with brucellosis. *Clin Vaccine Immunol* 2008 Feb;15(2):293-296.
- Shaheen BW, Nayak R, Foley SL, Kweon O, Deck J, Park M, et al. Molecular characterization of resistance to extended-spectrum cephalosporins in clinical *Escherichia coli* isolates from companion animals in the United States. *Antimicrob Agents Chemother* 2011 Dec;55(12):5666-5675.
- Alatoom A, Elsayed H, Lawlor K, AbdelWareth L, El-Lababidi R, Cardona L, et al. Comparison of antimicrobial activity between ceftolozane-tazobactam and ceftazidime-avibactam against multidrug-resistant isolates of *Escherichia coli*, *Klebsiella pneumoniae*, and *Pseudomonas aeruginosa*. *Int J Infect Dis* 2017 Sep;62:39-43.
- Sader HS, Flamm RK, Carvalhaes CG, Castanheira M. Comparison of ceftazidime-avibactam and ceftolozane-tazobactam *in vitro* activities when tested against gram-negative bacteria isolated from patients hospitalized with pneumonia in United States medical centers (2017-2018). *Diagn Microbiol Infect Dis* 2020 Mar;96(3):114833.
- Viala B, Zaidi FZ, Bastide M, Dumont Y, Le Moing V, Jean-Pierre H, et al. Assessment of the *in vitro* activities of ceftolozane/tazobactam and ceftazidime/avibactam in a collection of beta-lactam-resistant enterobacteriaceae and *Pseudomonas aeruginosa* clinical isolates at Montpellier University Hospital, France. *Microb Drug Resist* 2019 Nov;25(9):1325-1329.
- Araj GF, Berjawi DM, Musharrafieh U, El Beayni NK. Activity of ceftolozane/tazobactam against commonly encountered antimicrobial resistant Gram-negative bacteria in Lebanon. *J Infect Dev Ctries* 2020 Jun;14(6):559-564.
- Hirsch EB, Brigman HV, Zucchi PC, Chen A, Anderson JC, Eliopoulos GM, et al. Ceftolozane-tazobactam and

- ceftazidime-avibactam activity against  $\beta$ -lactam-resistant *Pseudomonas aeruginosa* and extended-spectrum  $\beta$ -lactamase-producing Enterobacterales clinical isolates from U.S. medical centres. *J Glob Antimicrob Resist* 2020 Sep;22:689-694.
22. Ortiz de la Rosa J-M, Nordmann P, Poirel L. ESBLs and resistance to ceftazidime/avibactam and ceftolozane/tazobactam combinations in *Escherichia coli* and *Pseudomonas aeruginosa*. *J Antimicrob Chemother* 2019 Jul;74(7):1934-1939.
  23. Bassetti M, Poulakou G, Timsit J-F. Focus on antimicrobial use in the era of increasing antimicrobial resistance in ICU. *Intensive Care Med* 2016 Jun;42(6):955-958.
  24. MacVane SH. Antimicrobial resistance in the intensive care unit: a focus on gram-negative bacterial infections. *J Intensive Care Med* 2017 Jan;32(1):25-37.
  25. Harris PN, Peleg AY, Iredell J, Ingram PR, Miyakis S, Stewardson AJ, et al. Meropenem versus piperacillin-tazobactam for definitive treatment of bloodstream infections due to ceftriaxone non-susceptible *Escherichia coli* and *Klebsiella* spp (the MERINO trial): study protocol for a randomised controlled trial. *Trials* 2015 Jan;16:24.
  26. Livermore DM, Mushtaq S, Meunier D, Hopkins KL, Hill R, Adkin R, et al; BSAC Resistance surveillance Standing Committee. Activity of ceftolozane/tazobactam against surveillance and 'problem' Enterobacteriaceae, *Pseudomonas aeruginosa* and non-fermenters from the British Isles. *J Antimicrob Chemother* 2017 Aug;72(8):2278-2289.
  27. Khan FY, Elshafie SS, Almaslamani M, Abu-Khattab M, El Hiday AH, Errayes M, et al. Epidemiology of bacteraemia in Hamad general hospital, Qatar: a one year hospital-based study. *Travel Med Infect Dis* 2010 Nov;8(6):377-387.
  28. Shaikh N, Momin U, Atef Shible A, Al-Musalmani M, Ansari A. Community acquired urosepsis: a surgical intensive care experience. *Qatar Med J* 2020 Apr;2020(1):8.
  29. Balkhy HH, Assiri AM, Mousa HA, Al-Abri SS, Al-Katheeri H, Alansari H, et al; at the workshop. The strategic plan for combating antimicrobial resistance in Gulf Cooperation Council States. *J Infect Public Health* 2016 Jul-Aug;9(4):375-385.
  30. Aly M, Balkhy HH. The prevalence of antimicrobial resistance in clinical isolates from Gulf Corporation Council countries. *Antimicrob Resist Infect Control* 2012 Jul;1(1):26.
  31. Khanfar H. Molecular epidemiological study of extended spectrum beta-lactamase (ESBL) producing bacteria from a hospital within Saudi Arabia. University of Portsmouth, 2017.
  32. Perez-Lopez A, Sundararaju S, Al-Mana H, Tsui KM, Hasan MR, Suleiman M, et al. Molecular characterization of extended-spectrum  $\beta$ -lactamase-producing *Escherichia coli* and *Klebsiella pneumoniae* among the pediatric population in Qatar. *Front Microbiol* 2020 Nov;11:581711.
  33. Abid FB, Tsui CK, Doi Y, Deshmukh A, McElheny CL, Bachman WC, et al. Molecular characterization of clinical carbapenem-resistant Enterobacterales from Qatar. *Eur J Clin Microbiol Infect Dis* 2021 Aug;40(8):1779-1785.
  34. Zalas-Więcek P, Gospodarek-Komkowska E. Antimicrobial susceptibility of multi-drug and extensively-drug-resistant *Escherichia coli* to ceftolozane-tazobactam and ceftazidime-avibactam: an in vitro study. *Postepy Hig Med Dosw* 2020;74:77-83.
  35. Ortiz de la Rosa J-M, Nordmann P, Poirel L. ESBLs and resistance to ceftazidime/avibactam and ceftolozane/tazobactam combinations in *Escherichia coli* and *pseudomonas aeruginosa*. *J Antimicrob Chemother* 2019 Jul;74(7):1934-1939.
  36. Lahiri SD, Alm RA. Identification of novel VEB  $\beta$ -lactamase enzymes and their impact on avibactam inhibition. *Antimicrob Agents Chemother* 2016 Apr;60(5):3183-3186.
  37. Barnes MD, Taracila MA, Rutter JD, Bethel CR, Galdadas I, Hujer AM, et al. Deciphering the evolution of cephalosporin resistance to ceftolozane-tazobactam in *pseudomonas aeruginosa*. *mBio* 2018 Dec;9(6):e02085-18.
  38. Arca-Suárez J, Lasarte-Monterrubio C, Rodiño-Janeiro B-K, Cabot G, Vázquez-Ucha JC, Rodríguez-Iglesias M, et al. Molecular mechanisms driving the in vivo development of OXA-10-mediated resistance to ceftolozane/tazobactam and ceftazidime/avibactam during treatment of XDR *Pseudomonas aeruginosa* infections. *J Antimicrob Chemother* 2021 Jan;76(1):91-100.
  39. Cultrera R, Libanore M, Barozzi A, d'Anchera E, Romanini L, Fabbian F, et al. Ceftolozane/tazobactam and ceftazidime/avibactam for multidrug-resistant gram-negative infections in immunocompetent patients: a single-center retrospective study. *Antibiotics (Basel)* 2020 Sep;9(10):640.
  40. Evans SR, Tran TT, Hujer AM, Hill CB, Hujer KM, Mediavilla JR, et al; Antibacterial resistance leadership group (ARLG). Rapid molecular diagnostics to inform empiric use of ceftazidime/avibactam and ceftolozane/tazobactam against *pseudomonas aeruginosa*: PRIMERS IV. *Clin Infect Dis* 2019 May;68(11):1823-1830.



**Table A1:** Microbiological characteristics, molecular characterization, and susceptibility testing results for 109 ESBL-producing Enterobacteriaceae isolates.

Isolate number	Collection mmm-yy	Organism	Location	Specimen type	Disk confirmation test	Molecular results			Antimicrobial susceptibility test (MIC)	
						SHV	TEM	CTXM1	CZA	C/T
1	Nov-12	<i>Klebsiella pneumoniae</i> ssp <i>pneumoniae</i>	SICU	Sputum	Positive	Negative	Negative	Positive	0.25	0.75
2	Nov-12	<i>Klebsiella pneumoniae</i> ssp <i>pneumoniae</i>	SICU	Urine	Positive	Positive	Positive	Positive	0.25	0.50
3	Nov-12	<i>Klebsiella pneumoniae</i> ssp <i>pneumoniae</i>	SICU	Sputum	Positive	Positive	Positive	Positive	0.75	1.00
4	Nov-12	<i>Escherichia coli</i>	SICU	Wound Swab	Positive	Negative	Negative	Positive	0.12	0.38
5	Nov-12	<i>Serratia marcescens</i>	MICU	Blood	Positive	Negative	Negative	Negative	0.12	0.19
6	Nov-12	<i>Klebsiella pneumoniae</i> ssp <i>pneumoniae</i>	PICU	Endotracheal Tube Secretion	Positive	Positive	Negative	Negative	0.19	0.38
7	Nov-12	<i>Klebsiella pneumoniae</i> ssp <i>pneumoniae</i>	NICU	Peritoneal fluid	Negative +AmpC	Negative	Positive	Positive	0.19	0.25
8	Nov-12	<i>Escherichia coli</i>	MICU	Urine	Negative +AmpC	Negative	Positive	Negative	0.50	2.00
9	Nov-12	<i>Klebsiella pneumoniae</i> ssp <i>pneumoniae</i>	SICU	Blood	Positive	Positive	Negative	Negative	0.75	1.50
10	Nov-12	<i>Klebsiella pneumoniae</i> ssp <i>pneumoniae</i>	PICU	Endotracheal Tube Secretion	Positive	Positive	Positive	Positive	0.19	0.38
11	Dec-12	<i>Klebsiella pneumoniae</i> ssp <i>pneumoniae</i>	SICU	Endotracheal Tube Secretion	Positive	Positive	Positive	Positive	0.19	0.38
12	Dec-12	<i>Klebsiella oxytoca</i>	NICU	Tracheostomy Site Swab	Positive	Negative	Negative	Negative	0.12	0.38
13	Dec-12	<i>Klebsiella pneumoniae</i> ssp <i>pneumoniae</i>	PICU	Urine	Positive	Positive	Positive	Positive	0.25	0.50
14	Dec-12	<i>Escherichia coli</i>	NICU	Conjunctival Swab	Positive	Negative	Positive	Positive	0.09	0.38
15	Dec-12	<i>Enterobacter aerogenes</i>	MICU	Blood	Positive	Positive	Negative	Negative	0.19	0.50
16	Dec-12	<i>Klebsiella pneumoniae</i> ssp <i>pneumoniae</i>	MICU	Endotracheal Tube Secretion	Positive	Positive	Positive	Positive	0.19	0.38
17	Jan-13	<i>Escherichia coli</i>	MICU	Urine	Negative	Negative	Negative	Negative	0.38	0.50
18	Jan-13	<i>Escherichia coli</i>	SICU	Blood	Positive	Negative	Positive	Negative	0.25	0.38
19	Jan-13	<i>Escherichia coli</i>	TICU	Blood	Positive	Negative	Positive	Positive	0.125	0.25

**Table A1:** Microbiological characteristics, molecular characterization, and susceptibility testing results for 109 ESBL-producing Enterobacteriaceae isolates.

-continued

Isolate number	Collection mmm-yy	Organism	Location	Specimen type	Disk confirmation test	Molecular results			Antimicrobial susceptibility test (MIC)	
						SHV	TEM	CTXMI	CZA	C/T
20	Jan-13	<i>Enterobacter cloacae</i>	TICU	Blood	Negative	Negative	Negative	Negative	0.12	0.25
21	Jan-13	<i>Klebsiella pneumoniae</i> sp <i>pneumoniae</i>	SICU	Sputum	Positive	Positive	Positive	Positive	0.25	0.38
22	Jan-13	<i>Escherichia coli</i>	MICU	Blood	Negative +AmpC	Negative	Negative	Negative	0.38	1.50
23	Jan-13	<i>Proteus penneri</i>	MICU	Blood	Positive	Negative	Negative	Negative	0.047	1.00
24	Jan-13	<i>Escherichia coli</i>	PICU	Urine	Positive	Negative	Negative	Positive	0.12	0.38
25	Jan-13	<i>Klebsiella pneumoniae</i> sp <i>pneumoniae</i>	MICU	Endotracheal Tube Secretion	Positive	Positive	Negative	Negative	0.25	0.38
26	Jan-13	<i>Klebsiella pneumoniae</i> sp <i>pneumoniae</i>	TICU	Urine	Positive	Positive	Positive	Positive	0.25	0.75
27	Jan-13	<i>Citrobacter braakii</i>	TICU	Blood	Negative	Negative	Negative	Negative	0.25	0.75
28	Jan-13	<i>Escherichia coli</i>	TICU	Sputum	Positive	Negative	Negative	Positive	0.94	0.38
29	Jan-13	<i>Escherichia coli</i>	PICU	Urine	Positive	Negative	Positive	Positive	0.12	0.38
30	Feb-13	<i>Serratia marcescens</i>	MICU	Sputum	Negative	Negative	Negative	Negative	0.02	0.19
31	Feb-13	<i>Escherichia coli</i>	MICU	Urine	Positive	Positive	Positive	Positive	0.06	0.25
32	Feb-13	<i>Escherichia coli</i>	MICU	Blood	Positive	Negative	Negative	Positive	0.12	0.50
33	Feb-13	<i>Klebsiella pneumoniae</i> sp <i>pneumoniae</i>	MICU	Tracheal Aspirate	Positive	Positive	Positive	Positive	0.19	0.38
34	Feb-13	<i>Klebsiella pneumoniae</i> sp <i>pneumoniae</i>	SICU	Urine	Positive	Positive	Negative	Positive	0.09	0.38
35	Feb-13	<i>Escherichia coli</i>	MICU	Blood	Positive	Negative	Negative	Positive	0.19	0.38
36	Mar-13	<i>Escherichia coli</i>	TICU	Ascitic Fluid	Positive	Negative	Negative	Positive	0.12	0.38
37	Mar-13	<i>Klebsiella pneumoniae</i> sp <i>pneumoniae</i>	TICU	Sputum	Positive	Positive	Negative	Negative	0.25	0.38
38	Mar-13	<i>Klebsiella pneumoniae</i> sp <i>pneumoniae</i>	SICU	Urine	Positive	Negative	Negative	Positive	0.38	0.38
39	Mar-13	<i>Klebsiella pneumoniae</i> sp <i>pneumoniae</i>	NICU	Blood	Positive	Positive	Negative	Positive	0.25	0.38
40	Mar-13	<i>Klebsiella pneumoniae</i> sp <i>pneumoniae</i>	SICU	Blood	Positive	Positive	Negative	Positive	0.25	0.38
41	Mar-13	<i>Escherichia coli</i>	SICU	Peritoneal fluid	Positive + AmpC	Negative	Negative	Positive	256.00	256.00

**Table A1:** Microbiological characteristics, molecular characterization, and susceptibility testing results for 109 ESBL-producing Enterobacteriaceae isolates.

-continued

Isolate number	Collection mmm-yy	Organism	Location	Specimen type	Disk confirmation test	Molecular results	Antimicrobial susceptibility test (MIC)
						SHV TEM CTXMI	CZA C/T
42	Mar-13	<i>Escherichia coli</i>	SICU	Peritoneal fluid	Negative + AmpC	Negative	0.19 0.75
43	Mar-13	<i>Escherichia coli</i>	SICU	Urine	Positive	Negative	0.12 0.38
44	Mar-13	<i>Klebsiella pneumoniae ssp pneumoniae</i>	PICU	Urine	Positive	Positive	0.19 0.5
45	Mar-13	<i>Klebsiella pneumoniae ssp pneumoniae</i>	MICU	Sputum	Positive	Positive	0.38 1.00
46	Mar-13	<i>Klebsiella pneumoniae ssp ozaenae</i>	MICU	Sputum	Positive	Negative	0.09 0.25
47	Mar-13	<i>Klebsiella pneumoniae ssp pneumoniae</i>	MICU	Tracheal Aspirate	Positive	Positive	0.12 0.25
48	Mar-13	<i>Klebsiella pneumoniae ssp pneumoniae</i>	MICU	Urine	Positive	Positive	0.38 1.50
49	Mar-13	<i>Escherichia coli</i>	TICU	Urine	Positive	Positive	0.09 0.38
50	Apr-13	<i>Klebsiella pneumoniae ssp pneumoniae</i>	TICU	J VAC Fluid	Positive	Positive	0.19 0.5
51	Apr-13	<i>Escherichia coli</i>	TICU	J VAC Fluid	Positive	Negative	0.09 0.75
52	Apr-13	<i>Escherichia coli</i>	PICU	Urine	Positive	Negative	0.12 0.38
53	Apr-13	<i>Klebsiella pneumoniae ssp pneumoniae</i>	PICU	Endotracheal Tube Secretion	Positive	Positive	0.19 0.75
54	Apr-13	<i>Enterobacter aerogenes</i>	MICU	Tracheal Aspirate	Positive	Positive	0.25 0.38
55	Apr-13	<i>Klebsiella pneumoniae ssp pneumoniae</i>	SICU	Blood	Positive	Negative	0.75 1.00
56	Apr-13	<i>Klebsiella pneumoniae ssp pneumoniae</i>	PICU	Endotracheal Tube Secretion	Positive	Negative	0.25 0.38
57	Apr-13	<i>Klebsiella pneumoniae ssp pneumoniae</i>	SICU	Blood	Positive	Negative	0.19 0.38
58	Apr-13	<i>Escherichia coli</i>	SICU	Wound Swab	Positive	Negative	0.12 0.50
59	Apr-13	<i>Klebsiella pneumoniae ssp pneumoniae</i>	SICU	Sputum	Positive	Negative	0.25 0.38
60	Apr-13	<i>Klebsiella pneumoniae ssp pneumoniae</i>	SICU	Blood	Positive	Negative	0.19 0.25
61	Apr-13	<i>Klebsiella pneumoniae ssp pneumoniae</i>	NICU	Blood	Positive	Positive	0.19 0.38

**Table A1:** Microbiological characteristics, molecular characterization, and susceptibility testing results for 109 ESBL-producing Enterobacteriaceae isolates.

-continued

Isolate number	Collection mmm-yy	Organism	Location	Specimen type	Disk confirmation test	SHV	TEM	CTXMI	CZA	C/T
62	Apr-13	<i>Klebsiella pneumoniae</i> ssp <i>pneumoniae</i>	MICU	Endotracheal Tube Secretion	Positive	Positive	Negative	Negative	0.25	0.25
63	May-13	<i>Citrobacter freundii</i>	TICU	Blood	Negative	Negative	Negative	Negative	0.06	0.38
64	May-13	<i>Escherichia coli</i>	MICU	Urine	Positive	Negative	Negative	Positive	0.06	0.25
65	May-13	<i>Klebsiella pneumoniae</i> ssp <i>pneumoniae</i>	PICU	Tracheostomy Site Swab	Positive	Positive	Positive	Positive	0.25	0.75
66	May-13	<i>Klebsiella pneumoniae</i> ssp <i>pneumoniae</i>	NICU	Endotracheal Tube Secretion	Positive	Positive	Positive	Positive	0.25	0.38
67	May-13	<i>Klebsiella pneumoniae</i> ssp <i>pneumoniae</i>	SICU	Sputum	Positive	Positive	Negative	Negative	0.19	0.38
68	May-13	<i>Klebsiella pneumoniae</i> ssp <i>pneumoniae</i>	TICU	Blood	Positive	Positive	Negative	Positive	0.19	0.75
69	May-13	<i>Klebsiella pneumoniae</i> ssp <i>pneumoniae</i>	MICU	Sputum	Positive	Positive	Negative	Negative	0.38	0.38
70	May-13	<i>Enterobacter cloacae</i>	SICU	Sputum	Positive	Positive	Negative	Negative	0.19	0.19
71	Jun-13	<i>Klebsiella pneumoniae</i> ssp <i>pneumoniae</i>	TICU	Wound Swab	Positive	Positive	Negative	Positive	0.38	0.38
72	Jun-13	<i>Klebsiella pneumoniae</i> ssp <i>pneumoniae</i>	SICU	Blood	Positive	Positive	Negative	Negative	0.19	0.38
73	Jun-13	<i>Klebsiella pneumoniae</i> ssp <i>pneumoniae</i>	SICU	Blood	Positive	Positive	Positive	Positive	0.19	1.00
74	Jun-13	<i>Escherichia coli</i>	PICU	Endotracheal Tube Secretion	Positive	Negative	Negative	Positive	0.12	0.38
75	Jun-13	<i>Escherichia coli</i>	PICU	Urine	Positive	Negative	Negative	Positive	0.12	0.38
76	Jun-13	<i>Klebsiella pneumoniae</i> ssp <i>pneumoniae</i>	SICU	Blood	Positive	Positive	Positive	Positive	0.38	1.50
77	Jul-13	<i>Klebsiella pneumoniae</i> ssp <i>pneumoniae</i>	NICU	Blood	Positive	Positive	Negative	Negative	0.38	0.38
78	Jul-13	<i>Escherichia coli</i>	NICU	Blood	Positive	Negative	Negative	Negative	0.12	0.38
79	Jul-13	<i>Klebsiella pneumoniae</i> ssp <i>pneumoniae</i>	MICU	Sputum	Positive	Positive	Negative	Positive	0.25	0.75
80	Jul-13	<i>Klebsiella pneumoniae</i> ssp <i>pneumoniae</i>	SICU	Endotracheal Tube Secretion	Positive	Positive	Positive	Positive	0.19	0.38
81	Jul-13	<i>Escherichia coli</i>	PICU	Urine	Positive	Negative	Negative	Positive	0.12	0.38



**Table A1:** Microbiological characteristics, molecular characterization, and susceptibility testing results for 109 ESBL-producing Enterobacteriaceae isolates.

-continued

Isolate number	Collection mmm-yy	Organism	Location	Specimen type	Disk confirmation test	SHV	TEM	CTXMI	CZA	C/T
82	Jul-13	<i>Escherichia coli</i>	SICU	Urine	Positive	Negative	Positive	Negative	0.25	0.50
83	Jul-13	<i>Klebsiella pneumoniae ssp pneumoniae</i>	MICU	BAL	Positive	Positive	Negative	Positive	0.25	0.75
84	Jul-13	<i>Escherichia coli</i>	MICU	Urine	Positive	Negative	Negative	Negative	0.09	0.25
85	Jul-13	<i>Citrobacter amalonaticus</i>	PICU	Urine	Positive	Positive	Negative	Negative	0.12	0.19
86	Jul-13	<i>Klebsiella pneumoniae ssp pneumoniae</i>	MICU	Sputum	Positive	Positive	Positive	Positive	0.25	0.50
87	Jul-13	<i>Enterobacter cloacae</i>	NICU	Eye Swab	Negative + AmpC	Negative	Negative	Negative	0.09	0.25
88	Jul-13	<i>Enterobacter cloacae</i>	TICU	Blood	Negative + AmpC	Negative	Negative	Negative	0.09	0.25
89	Jul-13	<i>Enterobacter aerogenes</i>	TICU	Blood	Positive	Positive	Negative	Negative	0.09	0.38
90	Aug-13	<i>Escherichia coli</i>	MICU	Ascitic fluid	Positive	Negative	Positive	Positive	0.12	0.25
91	Aug-13	<i>Klebsiella pneumoniae ssp pneumoniae</i>	MICU	Urine	Positive	Positive	Negative	Positive	0.38	0.75
92	Aug-13	<i>Enterobacter aerogenes</i>	TICU	Sputum	Positive	Positive	Negative	Negative	0.25	0.38
93	Aug-13	<i>Klebsiella pneumoniae ssp pneumoniae</i>	MICU	Sputum	Positive	Positive	Positive	Positive	0.19	0.75
94	Sep-13	<i>Escherichia coli</i>	NICU	Blood	Positive	Negative	Negative	Positive	0.19	0.75
95	Sep-13	<i>Klebsiella pneumoniae ssp pneumoniae</i>	PICU	Urine	Positive	Positive	Positive	Positive	0.25	0.5
96	Sep-13	<i>Escherichia coli</i>	MICU	Urine	Positive	Negative	Positive	Positive	0.09	0.38
97	Sep-13	<i>Klebsiella pneumoniae ssp pneumoniae</i>	MICU	Urine	Positive	Positive	Positive	Positive	0.25	0.75
98	Sep-13	<i>Escherichia coli</i>	SICU	Sputum	Positive	Negative	Negative	Positive	0.12	0.50
99	Sep-13	<i>Klebsiella pneumoniae ssp pneumoniae</i>	MICU	Sputum	Positive	Positive	Positive	Positive	0.25	0.75
100	Sep-13	<i>Klebsiella pneumoniae ssp pneumoniae</i>	PICU	Blood	Positive	Negative	Positive	Positive	0.12	0.25
101	Sep-13	<i>Escherichia coli</i>	PICU	Urine	Positive	Negative	Negative	Positive	0.06	0.25
102	Sep-13	<i>Klebsiella pneumoniae ssp pneumoniae</i>	MICU	Sputum	Positive	Positive	Positive	Positive	0.19	0.38
103	Sep-13	<i>Klebsiella pneumoniae ssp pneumoniae</i>	NICU	Central line Tip	Positive	Positive	Negative	Positive	0.25	0.50
104	Sep-13	<i>Escherichia coli</i>	TICU	Sputum	Positive	Negative	Positive	Positive	0.19	0.25
105	Oct-13	<i>Escherichia coli</i>	SICU	Blood	Positive	Negative	Negative	Positive	0.06	0.19

**Table A1:** Microbiological characteristics, molecular characterization, and susceptibility testing results for 109 ESBL-producing Enterobacteriaceae isolates.

-continued

Isolate number	Collection mmm-yy	Organism	Location	Specimen type	Disk confirmation test	Molecular results			Antimicrobial susceptibility test (MIC)	
						SHV	TEM	CTXMI	CZA	C/T
106	Oct-13	<i>Escherichia coli</i>	SICU	Sputum	Positive	Positive	Positive	Negative	0.06	0.19
107	Oct-13	<i>Escherichia coli</i>	SICU	Urine	Positive	Negative	Negative	Positive	0.09	0.25
108	Oct-13	<i>Klebsiella pneumoniae ssp pneumoniae</i>	SICU	Blood	Positive	Positive	Positive	Positive	0.38	1.00
109	Oct-13	<i>Escherichia coli</i>	TICU	Sputum	Positive	Positive	Positive	Negative	0.06	0.25

ESBL: extended-spectrum  $\beta$ -lactamase; MIC: minimum inhibitory concentration; White: susceptible; grey: non-susceptible, susceptibility was reported according to Clinical Laboratory Standards Institute (CLSI) breakpoints (Clinical Laboratory Standards Institute, 2020). m: month; y: year; MICU: Medical Intensive Care Unit; NICU: Neonatal Intensive Care Unit; PICU: Pediatric Intensive Care Unit; SICU: Surgical Intensive Care Unit; TICU: Trauma Intensive Care Unit; CZA: ceftazidime-avibactam; C/T: ceftolozane-tazobactam.