

# Effect of Two Different Doses of Dexmedetomidine as Adjuvant in Bupivacaine Induced Subarachnoid Block for Elective Abdominal Hysterectomy Operations: A Prospective, Double-blind, Randomized Controlled Study

---

ONLINE

DOI 10.5001/omj.2020.76

---

## **Retraction of the published manuscript:**

Das A, Halder S, Chattopadhyay S, Mandal P, Chhaule S, Banu R. Effect of Two Different Doses of Dexmedetomidine as Adjuvant in Bupivacaine Induced Subarachnoid Block for Elective Abdominal Hysterectomy Operations: A Prospective, Double-blind, Randomized Controlled Study. Oman Med J 2015 July; 30(4):257-263. DOI 10.50010/omj.2015.52.

The above article has been retracted due to self-plagiarism. Similarity of the abstract, table titles, and figure trends and caption with article titled Effect of Different Doses of Dexmedetomidine as Adjuvant in Bupivacaine -Induced Subarachnoid Block for Traumatized Lower Limb Orthopaedic Surgery: A Prospective, Double-Blinded and Randomized Controlled Study were noticed. Both articles have the same three authors. How to cite this retraction:

## **How to cite this retraction:**

Das A, Halder S, Chattopadhyay S, Mandal P, Chhaule S, Banu R. Effect of Two Different Doses of Dexmedetomidine as Adjuvant in Bupivacaine Induced Subarachnoid Block for Elective Abdominal Hysterectomy Operations: A Prospective, Double-blind, Randomized Controlled Study. Oman Med J 2015 July; 30(4):257-263. Retraction in: Oman Med J 2020 Jul:35(4):e155.