

External Ophthalmomyiasis due to *Oestrus Ovis*: A Case Report from Oman

Abdelhameed A. Abuelhassan

Abstract

Human myiasis has been reported from various regions in the world. We report a human case of external ophthalmomyiasis caused by the larvae of a sheep nasal botfly, *Oestrus ovis*, from Wadi Alhawasna, North Batina, Oman. A 9-years-old boy presented with symptoms of acute conjunctivitis. Five larvae were removed from his eye and sent for identification. The patient recovered completely after removal of the larvae.

From the Department of General Practitioners, Wadi Al Hawasna Hospital.

Received: 30 June 2010

Accepted: 14 Aug 2010

Address correspondence and reprint request to: Abdelhameed A. Abuelhassan, Medical Officer In-Charge, Wadi Al Hawasna Hospital, P.O. Box 440, Khaboura 326, North Batina, Sultanate of Oman.

Email: abdelhamid1427@yahoo.com

doi:10.5001/omj.2010.94

Introduction

Myiasis is the infestation of tissues or organs of animals or man by fly larvae. Human ophthalmomyiasis is mainly seen in people who are in contact with animals, in particular sheep and goats. The sheep nasal botfly, *Oestrus ovis*, is by far the commonest cause of human ophthalmomyiasis. The normal hosts for the larvae are sheep and goats. Occasionally, humans are the intermediate hosts.

Case Report

A 9-years-old boy presented to the Outpatient Department in February 2009 with a 1-day history of foreign body sensation and excessive watering from his right eye. He developed these symptoms after being struck in his right eye by a fly. He was absolutely fine before that and had no significant ocular or medical history. On examination; his visual acuity was 6/6 in both eyes. Examination of his left eye was normal. Eyelids of the affected right eye were normal. The conjunctiva was mildly congested with profuse lacrimation in the right eye. Five Larvae were removed with cotton buds from the conjunctiva, preserved in 70% alcohol and sent to the Veterinary Research Centre, Ministry of Agriculture and fisheries, Muscat for identification. Topical antihistaminic and antibiotic drops were given to the patient. The patient came for a follow-up after two days and he was completely relieved of his symptoms. Examination of both his eyes was normal. The larvae were identified as the first instar of *Oestrus Ovis*. (Fig. 1)

Discussion

Ophthalmomyiasis is classified as ophthalmomyiasis externa if the larvae are present in the conjunctiva and ophthalmomyiasis interna when there is intraocular penetration of larvae.¹ The symptoms of ophthalmomyiasis externa are of acute conjunctivitis with soreness,



Figure 1: First instar of *Oestrus ovis*

lacrimation and conjunctival congestion. The symptoms of rhinitis in association with conjunctival reaction have been reported.² External ophthalmomyiasis can be complicated by corneal ulcer, decreased vision or invasion into eye globe. Apart from mechanical removal, there is no other therapy described. Topical antibiotics are useful in preventing secondary infection by bacteria. Cases of ophthalmomyiasis externa have been reported from various parts of the world.^{1,3} To the best of our knowledge, this represents the third case reported from Oman of ophthalmomyiasis caused by *Oestrus Ovis*.^{4,5}

Conclusion

Doctors in Oman should suspect ophthalmomyiasis in unilateral conjunctivitis with a foreign body sensation.

Acknowledgements

The author reported no conflict of interest and no funding was received on this work.

References

1. Gupta VP, Baveja UK. Ophthalmomyiasis externa caused by the sheep nasal botfly *Oestrus ovis*. *Indian J Ophthalmol* January-March 1988;36(1):41-42.
2. Mazzeo V, Ercolani D, Trombetti D, Todeschini R, Gaiba G. External ophthalmomyiasis. Report of four cases. *Int Ophthalmol* December 1987;11(2):73-76.
3. McDermott S, Schafer MD. Ophthalmomyiasis due to *Oestrus ovis* in South Australia. *Med J Aust* February 1983;1(3):129-130.
4. Victor R, Bhargva K. Ophthalmomyiasis in Oman: a case report and comments. *Wilderness Environ Med* 1998;9(1):32-35.
5. Shenoy R, Yanamadala S, Idris MA. Follicular conjunctivitis secondary to external ophthalmomyiasis due to larvae of *oestrus ovis*, *Delhi journal of ophthalmology* 2001; 9(1):73-75.